

# Review An Investigation of The Effects of Macro and Micro elements on the Production of High-Quality Seed

S. Patra\*

Crop Research and Seed Multiplication farm-to be tenable at Botany Department The University of Burdwan, Burdwan-713104, East Burdwan, West Bengal, India

## ABSTRACT

*For optimal development, growth, and ultimately the production of quality seed in each and every crop, requirements of macro and microelements are critical. If crops are to be produced as seed crops then the seed crops must meet the same macro and micronutrient requirements. In this regard, a review was conducted to examine the effects of macro and micro elements on seed quality. According to the results of numerous studies, the macronutrients like nitrogen, phosphorus, potassium, sulphur, calcium, magnesium and micronutrients like boron, chloride, iron, manganese, molybdenum, zinc have a significant impact on seed quality, which in turn affects seed yield.*

**Keywords-** macroelements, microelements; seed germination, seed vigour

## INTRODUCTION

Seeds available in the market are often very poor in quality in respect of germination, varietal purity, seed viability etc. The scarcity of quality seeds of crops in India during growing season is important issue till now due to lower production of quality seeds in a limited area. The possible way to produce the maximum quantity of quality seed, through manipulation of existing method of cultivation such as planting geometry, fertilization, irrigation and other cultural management practices are essential. The quality of seeds plays a critical role in the economic production of all crops. In this connection, numerous articles detailing the impact of changes in seed quality on seedling emergence and crop growth in a variety of species have been published (Powell, Matthews, and Oliveria, 1984; Tekrony and Egli, 1991).

Poor crop stand can promote weed populations causing competition for water, light and limiting nutrients (Kropff and van Laar, 1993). It has been shown that the absence of some essential elements, such as Mg and Zn in most of the chemical fertilizers may be partly responsible for the low yield of maize in the semi-arid zones of northern Ghana (Abunyewa and Mercer-Quarshie 2004). Micronutrient deficiency is severing problem in soil and plants worldwide (Imtiaz et al., 2010) while appropriate quality of micronutrients is necessary for better growth, better flowering, higher fruit set, higher yield, quality and post-harvest life of horticultural products (Raja, E. M., 2009; Ram, R. A., Bose, T. K., 2000; Shekhar C et al, 2010; Sourour M.M., 2000) while its deficiency leads in lowering the productivity (Karuna, S. et al, 2019 and Zagade P.M et al., 2017). A staple fertilization program with macronutrients and micronutrients in plant nutrition is very essential in the high production of crop yield with good quality seed, so there is a need balance use of fertilizers and agronomic procedures are needed to increase yield of this crop. The function of macronutrients and micronutrients is vital in crop nutrition for improved yield and seed quality (Saeed B et al., 2012). The soil contains the majority of the nutrients that plants need to build their tissues, produce their nucleic acids, enzymes, and maintain their osmotic equilibrium. For healthy and productive crops, plants must

maintain a balance of 15 components, 12 of which are provided by the soil and are controlled by farmers (e.g., macro nutrients like nitrogen, phosphorus, potassium, sulfur, magnesium, calcium and micro nutrients like iron, boron, manganese, zinc, molybdenum, copper). Mineral nutrient stores in the seed must be sufficient to maintain seedling growth until root uptake starts to feed soil nutrients. Numerous studies have shown the significance of this, particularly for crops grown on soils with low nitrogen supply (Asher, 1987). One of the various methods used in seed farming to address seed nutrition at its most crucial point, i.e. seed filling, is supplemental foliar treatment (Shibles et al., 1975). It is generally known that major, minor, and macronutrients can be applied to seed crops through foliar spray (De, 1967; Kramer, 1969), and that micronutrients in particular, can improve the qualities of seeds more than soil application or dusting (Sabir Ahmed, 1989). Crop cultivation depends on a healthy soil environment. How well seeds establish, grow, and produce quality seeds is greatly influenced by the soil fertility, pH of the soil, microbial community, structural and textural quality of the soil, and other elements of the soil. There is a lack of information on this subject, and only a few studies, namely the impact of macro- and micronutrients on seed performance, have been thoroughly studied by academics. These are listed below:

## ARTICLE HISTORY

19 September 2022: Received  
17 January 2023: Revised  
28 March 2023: Accepted  
18 April 2023: Available Online

DOI:

<https://doi.org/10.5281/zenodo.8114785>

CORRESPONDING AUTHOR:

Sabyasachi Patra

E-MAIL ID:

sabya1983@gmail.com

COPYRIGHT:

© 2023 by the authors. The license of Theoretical Biology Forum. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

### **Effect of Nitrogen**

Application of nitrogen (N) encourages soil to absorb nutrients (Halder NK et al, 1998). When N application is increased, the oil quality deteriorates as a result of an increase in erucic acid and a drop in the concentrations of linoleic and oleic acids (Gupta et al., 2002). Okra (*Abelmoschus esculentus*) seeds from parent plants grown in fields with high nitrogen levels exhibited considerably better pre- and post-storage germination rates than seeds from plants that did not receive additional nitrogen fertilizer (Das 1990). According to Austin (1972), A significant development of germination inhibitors in the fruits of high-nitrogen plants may be the cause of the reduced geminability of high-nitrogen fruit observed in previous investigations (Sneddon, 1963; Scott, 1969).

### **Effect of Phosphorus**

Phosphorus is essential for the synthesis of ADP and ATP, phospholipids, and nucleic acids as well as the activity of enzymes (Thompson LR and Torch FR, 1978). Therefore, it is essential for the development of roots, flowers, fruits, and seeds (Smil 2000). Talooth et al., (1989) studied how different sources of Phosphorus affected soybean growth, and found that when applied at a rate of up to 150 kg per hectare, all Phosphorus fertilizers increased soybean seed yield. Raboy and Dickinson (1987) observed that increasing the total Phosphorus nutrition from the third week after flowering linearly increased its transfer to growing seeds till physiological maturity. When a soybean reaches physiological maturity, the seed quality is at its maximum (Egli and Tekrony, 1995) and if seed is not harvested at this stage, the seed quality declines due to daily fluctuation in the seed moisture and field environment.

### **Effect of potassium**

Studies on wheat have shown that a deficiency in potassium supply to the planting material reduces the development period of the developing grain, which is accompanied with an earlier presence of the ABA maximum and an enhancement in its exact amount. (Haeder & Beringer, 1981a). Potassium helps to increase the absorption of nitrogen (Sing SP, Verma AB. 2001; Vigidal SM). It has been implicated with the maintenance of good seed quality as significantly lower levels of diseased seed produced when levels of potassium in the soil increased (Crittenden and Svec, 1974).

### **Effect of calcium**

Burton et al. (2000) investigated the effects of Ca on soybean seed quality and production and discovered that lower Ca levels affected seed filling, seed production, seed germination, and the occurrence of aberrant seedlings. Similar negative consequences of low Ca levels in plants leading to poor seed quality issues such decreased germination and vigor have been documented (Keiser and Mullen, 1993). Loss of membrane integrity is the main consequence of calcium shortage in plant cells (Hetch-Buchholz, 1979) and a faulty and stunted root system is a sign of Ca-deficient plants (Nelson and Niedziela 1998); Therefore, it is crucial to externally provide nutrients for crops with shallow roots, like onions, that are unable to collect nutrients from the deep soil profile. Sharma et al. (2010) found that mineral nutrients should be supplied during an active growth stage in order to achieve rapid plant development, a healthy plant stand, a bigger seed harvest, and increased seed quality.

### **Effect of Magnesium**

It stimulates the most enzymes in comparison to other mineral nutrients (Epstein and Bloom 2005). Due to its role as the activator of vital photosynthetic enzymes and the core atom in the chlorophyll molecule, magnesium is a vital component of photosynthesis (Wedding and Black 1988; Portis 1992; Marschner 2012). Seed reserves are a major factor in germination and seedling emergence such as starch (Douglass et al. 1993; Wang et al. 2011), and consequently, for durum wheat, a clear relationship between seed size and seedling vigour was observed (Royo et al. 2006). Quick production and use of ATP is also necessary during germination (Perl 1986; Weitbrecht et al. 2011). The content of ATP and the ability of seeds to germinate were found to be highly correlated in a prior study (Siegenthaler and Douet-Orhant 1944). Mg is necessary for ATP's synthesis and operation (Boyer 1997; Marschner 2012; Igamberdiev and Kleczkowski 2015), A lack of magnesium in the seed may potentially hinder germination by altering the availability and physiological function of ATP. Mg levels must be enough for sugar transfer in germination of seeds and growth of seedlings. In order to maintain germination, sucrose is transported from the scutellum into the actively growing sink tissues of roots and shoots (Aoki et al. 2006; Scofield et al. 2007) and similarly, as it is in leaves, the translocation of sugars into the sink tissues of sprouting seeds is probably dependent on an adequate supply of magnesium. Chlorosis, caused by a magnesium deficit, causes the leaves to turn yellow and deteriorate the chlorophyll content; yet, a sufficient amount of magnesium makes the plant healthy (Hermans et al., 2010)

### **Effect of Boron**

Oilseed rape has been categorized as a crop that is extremely susceptible to B shortage, hence B nutrition is crucial to its growth. This seems to be caused in part by strong demand for this micronutrient (Camacho-Cristobal et al. 2008; Dell and Huang 1997; Herrera-Rodriguez et al. 2010; Stangoulis et al. 2000; Zhang et al. 2014). The growth of roots and shoots is generally slowed or inhibited by inadequate B supply to oilseed rape because it restricts cell expansion and cell division, affects the formation of vascular bundles, and lowers pollen germination and pollen tube growth. Consequently, B deficiency results in decreased seed quality (Dell and Huang 1997), seed abortion, damaged embryos, or shrivelled seeds, lower seed germination, and reduced seed vigour in the following generation of oilseed rape seedlings (Zhao et al. 2012). While seed priming with B (0.02% solution of H<sub>3</sub>BO<sub>3</sub>) had no noticeable impact on seed germination in oats (*Avena sativa* L.), it did enhance tillering, panicle length, and grain weight, which led to an 8.42% increase in grain production when compared to untreated seeds (Saric and Saciragic, 1969).

### **Effect of Molybdenum**

Oilseed rape, one of the most significant commercial and agricultural crops, is sensitive to Mo deficiency in soil, which affects its growth, yield, and seed quality. Liu et al. (2010a) showed that varied amounts of phosphorus (P) had an impact on how much dry matter, chlorophyll concentration, and net photosynthesis ratio (Pn) of seedling leaves changed with the addition of Mo to oilseed rape. Additionally, P fertiliser was clearly the only factor that boosted oilseed rape yield. (Liu et al. 2010b). Yang et al. (2009) also reported that the addition of zinc (Zn) and boron (B) fertilizers led to an increase in seed

production when Mo was present. However, with the application of Mo and B fertilizer, the seed quality, including the oil content and oleic and linoleic acid contents, increased greatly, whereas the quantities of stearic acid, sulfuric glucosid, and erucic acid declined (Chen et al. 2004). Similar results were also demonstrated under application of Mo and P fertilizers (Liu et al. 2012).

### Effect of Mn

Storing of mineral nutrient in seed play a significant role in determining vigour and standard seedling development. Acute Mn deficiency can cause dark necrotic patches to form on leaves, which can cause early leaf loss. Some cereal leaves have white and grey patches, which are symptoms of a Mn deficit (Chang, 1999; Stout and Arnon, 1939). Low manganese (Mn) content in seed resulted in wheat seedlings with low vigour and low yields (Marcar and Graham, 1986; Singh and Bharti, 1985). Barley and lupin with low Mn level in seed showed delayed germination and seedling development. (Genc et al., 2000; Longenecker et al., 1991; Crosbie et al., 1994). Compared to the untreated control, priming *Echinacea purpurea* (L.) seed with a 0.1% MnSO<sub>4</sub> solution improved germination by 36% and field emergence by 27% (Babaeva et al., 1999). In comparison to soil application, all of the research done on seed priming with Mn have shown a significant improvement in plant establishment, growth, yield, and grain enrichment (Babaeva et al., 1999; Khalid and Malik, 1982; Marcar and Graham, 1986).

### Effect of copper

Plants lacking in copper are more susceptible to disease, particularly ergot (a fungus causing reduced yield and grain quality; Solberg et al., 1999). The emergence of disease-related symptoms can make it difficult to distinguish them from symptoms of a copper deficiency. The most vulnerable wheat varieties to copper deficiency are winter and spring wheat (Solberg et al., 1999)

### Effect of Sulphur

The responses of different oilseed crops to sulphur application differed significantly (Misra et al., 2002). Due to its higher concentrations of sulphate and S-containing secondary metabolites as compared to other species like wheat, oilseed rape (*Brassica napus* L.) is a crop with a high S demand (Oenema and Postma, 2003). The pungency of onions is increased by sulphur fertilizer (Paterson DR, 1979). The growth and yield of oil seed crops are typically negatively affected by sulphur shortage as a result of inadequate nutrient supply (Basumatary et al. 2019). According to Rakesh et al. (2016), As a component of the amino acids methionine, cysteine, and cystine, which are necessary for the synthesis of other molecules like coenzyme-A, thiamine, and glutathione as well as for the production of chlorophyll, sulphur plays a crucial role in the growth and development of crops. As a result, plants require this element for oxidation-reduction processes, the production of chlorophyll, and the synthesis of proteins (Kumar and Trivedi, 2011 and Pandey and Ali, 2012).

### Effect of Fe and Zn

Iron is transported from the seed coat to the embryo by passing through membranes and the apoplast that separates the two tissues (Wolswinkel, 1992). There is a lack of knowledge on the

variables that influence how Fe moves into and out of the seed coat (Marentes and Grusak, 1998), and to make transgenic manipulation easier and to enhance seed [Fe] (Grusak, 2002).

Applying Fe and Zn may have improved photosynthates translocation to the sink, boosted vigour, and accumulated more photosynthates (Kanda and Dixit 1995). In calcareous soil and particularly so in high-pH, drought-prone areas, Fe and Zn shortages in rice frequently arise, which can severely inhibit initial growth and vigour and ultimately reduce rice yield (Katyal and Randhawa 1983). Inter-veinal chlorosis of younger leaves is a symptom of iron deficiency in rice plants, whereas apparent symptoms of zinc deficiency in rice include wilting caused by loss of turgidity, retarded plant development, basal chlorosis, bronzing of leaves, and in some cases seedling death (Neue et al. 1998). The antifungal properties of zinc and other mineral nutrients, as well as zinc's potential role in the establishment of disease resistance in plants, may be responsible for the decrease in disease incidence and severity under mineral nutrient spray (Graham and Webb 1991). Imran et al. (2014) found that nutrient seed priming increases maize seedling development under low root zone temperature during early growth as well as a large increase in seed contents of the relevant nutrients, i.e., Fe (25%), Zn (500%) Borax (0.50%) and ZnSO<sub>4</sub> (0.25%) seed soaking enhanced papaya (*Carica papaya* L.) stand establishment and growth in the field (Deb et al., 2010). Kaur et al. (2009) further found that seeds treated in Zn (ZnSO<sub>4</sub>) for 12 hours increased *Chlorophytum borivilianum* (L.) germination. Similarly, soaking seeds in solutions of Zn-EDTA and fritted Zn increased yield and Zn uptake, but the efficiency of each source's Zn uptake varied (Kang and Okoro, 1976).

## CONCLUSIONS

Having revised the overall picture on effect of macro and micronutrients on seed performance, it is clearly stated the impact and non impact portion at different stages of crop growth and development. Therefore, farmers and growers who are involved in seed production for harvesting quality seed, should be aware initially about the effect of macro and micronutrients for applying good management practices.

## ACKNOWLEDGEMENTS

Thanks to the staff of Crop Research and Seed Multiplication Farm, The University of Burdwan for collecting web based information

## CONFLICT OF INTEREST

No conflict of interest.

## REFERENCES

1. Abunyewa, A. A., and H. Mercer-Quarshie. 2004. Response of maize to magnesium and zinc application in the semi-arid zone of West Africa. *Asian Journal of Plant Sciences* 3(1):1-5. doi:10.3923/ajps.2004.1.5.
2. Aoki N, Scofield GN, Wang XD, Offler CE, Patrick JW, Furbank RT (2006) Pathway of sugar transport in germinating wheat seeds. *Plant Physiol* 141:1255-1263. doi:10.1104/pp.106.082719



3. Asher, C. J. 1987. Crop nutrition during establishment phase: Role of seed reserves. In: Crop Establishment Problem in Queensland Recognition Research and Resolution, eds. I. M. Wood, W. H. Hazard, and F. Rom, pp. 88–106. Brisbane: The Australian Institute of Agricultural Science and Technology.
4. Austin, R.B. Effect of environment before harvesting on viability. In *Viability of Seeds*, ed. E.H. Roberts (London: Chapman and Hall, 1972), pp. 114-149.
5. Babaeva, E.Y., Volobueva, V.F., Yagodin, B.A., Klimakhin, G.I. 1999. Sowing quality and productivity of *Echinacea purpurea* in relation to soaking the seed in manganese and zinc solutions. *Izvestiya Timiryazevskoi Sel'skokhozyaistvennoi Akademii*. 4, 73-80.
6. Basumatary, A, Goswami, K. Ozah, D. Hazarika, S. and Timsena G. (2019) Integrated sulphur management in rape seed (*Brassica Canipestris*) Blackgram (*Vigna mungo*) sequence in Inceptisol of Assam. *Annals of Plant and Soil Research* 21(1): 7-13.
7. Boyer PD (1997) The ATP synthase - a splendid molecular machine. *Annu Rev Biochem* 66:717-749. doi: 10.1146/annurev.biochem.66.1.717
8. Burton, M. G., M. J. Laver, and M. B. McDonald. 2000. Calcium effects on soybean seed production, Elemental concentration, and seed quality. *Crop Sci*. 40: 476–482.
9. Camacho-Cristobal JJ, Rexach J, Gonzalez-Fontes A (2008) Boron in plants: deficiency and toxicity. *J Integr Plant Biol* 50:1247-1255. doi:10.1111/j.1744-7909.2008.00742
10. Chang, S.S. 1999. Micronutrients in crop production of Taiwan. In: *Proceedings of International Workshop on Micronutrient in Crop Production*, held Nov. 8-13, 1999, National Taiwan University, Taipei, Taiwan ROC.
11. Chen G, Nian F, Wang Y, Xu F 2004: Effect of B, Mo on fatty acid component of *Brassica napus*. *Chinese J. Oil Crop Sci.*, 26(2), 69–71. ((in Chinese).
12. Crosbie, J., N. E. Longnecker, and A. D. Robson. 1994. Seed manganese affects the early growth of lupins in manganese deficient conditions. *Australian Journal of Agricultural Research* 45: 1469–1482.
13. Das, C.S. Quality seed production and seed invigoration of okra (*Abelmoschus esculentus* L.). Ph.D. dissertation, University of Calcutta, 1990.
14. De, R. (1967). A review of foliar applications to crops. *Ind. Fert. News.*, 16 (12): 77-81.
15. Dell B, Huang LB (1997) Physiological response of plants to low boron. *Plant Soil* 193:103-120. doi:10.1023/ 1004264009230
16. Douglass SK, Juvik JA, Splittstoesser WE (1993) Sweet corn seedling emergence and variation in kernel carbohydrate reserves. *Seed Sci Technol* 21:433^445
17. Egli, D. B., and TeKrony, D. M. 1995. Soybean seed germination, vigor and field emergence. *Seed Sci. and Technology* 23: 595–607.
18. Epstein E, Bloom AJ (2005) *Mineral nutrition of plants: Principles and perspectives*, 2nd edn. Sinauer Associates, Sunderland
19. Genc, Y, G. K. McDonald, and R. D. Graham. 2000. Effect of seed zinc content on early growth of barley (*Hordeum vulgare* L.) under low and adequate soil zinc supply. *Australian Journal of Agricultural Research* 51: 37–46. Gericke, S., and B. Kurmi
20. Grusak M A 2002 Enhancing mineral content in plant food products. *J. Am. Coll. Nutr.* 21, 178S-183S.
21. Graham, R. D., and M. J. Webb. 1991. Micronutrients and disease resistance and tolerance in plants. *Micronutrients in agriculture, (micronutrients i2)* 4:329–370.
22. Gupta, S.K., Chaudhary, M.L., Das, S.N.(2002). Effect of nickel and nitrogen on seed yield, oil content and oil quality of raya (*Brassica juncea* L.) in typic ustochrept and typic haplaquept of Haryana. *Journal of Indian Society of Soil Science*. 50(1), 144-146.
23. Halder NK, Zaman MWU, Chowdhury MMU, Ullah MH, Nag BL. Effect of nitrogen and phosphorus on the uptake of different nutrient elements by onion bulb. *Bangladesh J. Sci. Indust. Res.* 1998;33(3):404-408.
24. Haeder HE, Beringer H. 1981a. Influence of potassium nutrition and water stress on the content of abscisic acid in grains and flag lea\ -es of wheat during grain development. *Journal of the Science of Food and Agriculture* 32: 552-556.
25. Herrera-Rodriguez MB, Gonzalez-Fontes A, Rexach J, Camacho- Cristobal JJ, Maldonado JM, Navarro-Gochicoa MT (2010) Role of boron in vascular plants and response mechanisms to boron stresses. *Global Science Books* 4:115-122
26. Hermans C, Vuylsteke M, Coppens F, Cristescu SM, Harren FJ, Inzé D et al. Systems analysis of the responses to long-term magnesium deficiency and restoration in *Arabidopsis thaliana*. *New Phytol*, 2010; 187(1):132- 144.
27. Igamberdiev AU, Kleczkowski LA (2015) Optimization of ATP synthase function in mitochondria and chloroplasts via the adenylate kinase equilibrium. *Front Plant Sci* 6:10. doi: 10.3389/fpls.2015.0001

28. Imran S, Afzal S, Amjad, M, Akram A, Khawar K M and Pretorius S (2014) Seed priming with aqueous plant extracts improved seed germination and seedling growth under chilling stress in lentil (*Lens culineris* Medik). *Acta Adv. in Agril Sci*, 2:58- 59.
29. Kanda, C., and L. Dixit. 1995. Effect of zinc and nitrogen fertilization on summer rice. *Indian Journal of Agronomy* 40 (2):695-97.
30. Katyal, J. C., and N. S. Randhawa. 1983. *Micronutrients*. FAO Fertilizer and Plant Nutrition Bull.7. Rome, Italy: FAO. Kumar, R. 2001. Effect of potassium and iron on growth, yield and nutrient uptake of rice. Ph.D. Thesis, submitted to Department of Agronomy, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi, U P, (India).
31. Kang BT, Okoro EG. Response of flooded rice grown on a vertisol from northern Nigeria to zinc sources and methods of application. *Plant Soil*. 1976;44(1):15-25. doi: 10.1007/BF00016951.
32. Karuna, S., Misra, S., Topwal, M., Singh, V.K., (2019). A Research Review on Use of Micronutrient in Fruit Crops. *Int.J.Curr.Microbiol.App.Sci*. 8(08), 3014-3025. doi: <https://doi.org/10.20546/ijcmas.2019.808.349>
33. Keiser, J. R., and R. E. Mullen. 1993. Calcium and relative humidity effects on soybean seed nutrition and seed quality. *Crop Sci*. 33: 1345-1349.
34. Khalid, B.Y., Malik, N.S.A. 1982. Presowing soaking of wheat seeds in copper and manganese solutions. *Commun. Soil Sci. Plant Anal.* 13, 981-986.
35. Kramer, P.J. (1969). *Plant and Soil water relationship- a modern synthesis*. TMH Edition. New delhi. Pp.253-254.
36. Kropff, M. J., and H. H. van Laar. 1993. *Modeling Crop-Weed Interactions*. Wallingford, UK: CAB International.
37. Liu H, Hu C, Hu X, Nie Z, Sun X, Tan Q, Hu H 2010a: Interaction of molybdenum and phosphorus supply on uptake and translocation of phosphorus and molybdenum by *Brassica napus*. *J. Plant Nutr.*, 33, 1751-1760. doi:10.1080/01904167.2010.503778
38. Liu H, Hu C, Sun X, Tan Q, Nie Z, Hu X 2010b: Interactive effects of molybdenum and phosphorus fertilizers on photosynthetic characteristics of seedlings and grain yield of *Brassica napus*. *Plant Soil*, 326, 345- 353. doi:10.1007/s11104-012-1289-1
39. Liu H, Hu C, Nie Z, Sun X, Tan Q 2012: Interactive effects of molybdenum and phosphorus fertilizers on grain yield and quality of *Brassica napus*. *Plant Nutr. Fertil. Sci.*, 18, 678-688. (in Chinese).
40. Longenecker, N. E., N. E. Marcar, and R. D. Graham. 1991. Increased manganese content of barley seeds can increase grain yield in manganese-deficient conditions. *Australian Journal of Agricultural Research* 42: 1065-1074.
41. Marentes E and Grusak M A 1998 Iron transport and storage within the seed coat and embryo of developing seeds of pea (*Pisum sativum* L.). *Seed Sei. Res.* 8, 367-375
42. Marcar, N. E., and R. D. Graham. 1986. Effect of seed manganese content on the growth of wheat (*Triticum aestivum*) under manganese deficiency. *Plant and Soil* 96: 165-173.
43. Marschner P (2012) *Marschner 's mineral nutrition of higher plants*, 3rd edn. Elsevier, Academic press, USA.
44. Mishra, S.K., Singh, R.N., Tiwari, V.N. and Tiwari, K.N. (2002) Effect of sulphur fertilization on quality characteristics of mustard seed in five cultivars. *Journal of Indian Society of Soil Science* 50: 141-142.
45. Neue, H. U., C. Quijano., D. Senadhira., and T. Setter. 1998. Strategies for dealing with micronutrient disorders and salinity in lowland rice systems. *Field Crops Research* 56:139-55. doi:10.1016/S0378-4290(97)00125-1.
46. Nelson, P. V., and C. E. Niedziela. 1998. Effects of calcium source and temperature regime on calcium deficiency during hydroponic forcing of tulip. *Scientia horticulturae* 73 (2):137-150. doi:10.1016/S0304-4238(97)00157-X.
47. Oenema, O., and Postma, R. (2003). "Managing sulphur in agroecosystems" in *Sulphur in Plants*, eds Y. P. Abrol and A. Ahmad (Dordrecht: Springer), 45-70. doi: 10.1007/978-94-017-0289-8\_3
48. Perl M (1986) ATP synthesis and utilization in the early stage of seed germination in relation to seed dormancy and quality. *Physiol Plant* 66:177-182. doi: 1 0. 1 1 1 1 /j. 1 399-3054. 1 986. tb01253.x
49. Paterson DR. Sulphur fertilization effect on yield and pungency. *Progress Report NO. PR-3551*, Texas Agril. Exp. Stat. 1979;2.
50. Powell AA, Matthews S, Oliveira MA. 1984. Seed quality in grain legumes. *Advances in Applied Biology* 10, 217-285
51. Portis AR Jr (1992) Regulation of ribulose 15-bisphosphate carboxylase/oxygenase activity. *Annu Rev Plant Physiol* 43:415-437. doi: 1 0. 1 146/annurev.pp.43.060 192.0022
52. Rakesh, S. and Banik, Ganesh Chandra (2016) Effect of sulphur levels and sources on growth, yield and quality of mustard in terai region of West Bengal. *Annals of Plant and Soil Research* 18(2): 152-155.

53. Raboy, V., and D. B. Dickinson. 1987. The timing and rate of phytic acid accumulation in developing soybean seeds. *Plant Physiol.* 85: 841-844.
54. Raja, E. M., (2009). Importance of micronutrients in the changing horticultural scenario. *J. Hort. Sci.*, 4 (1), 1-27.
55. Ram, R. A., Bose, T. K., (2000). Effect of foliar application of magnesium and micronutrients on growth, yield and fruit quality of mandarin orange (*Citrus reticulata* Blanco). *Indian Journal of Horticulture*, 57(3), 215-220.
56. Royo C, Ramdani A, Moragues M, Villegas D (2006) Durum wheat under mediterranean conditions as affected by seed size. *J Agron Crop Sci* 192:257-266. doi: 10.1111/j.1439-037X.2006.00215.x
57. R. Kaur, S. Arora and A.K. Thukral, 2009. Enhancing Seed Germination of *Chlorophytum borivillianum* Sant. Et Fernand. with PGRs, Steroidal Hormones and Zinc. *Research Journal of Seed Science*, 2: 32-39.
58. Saric, T., Saciragic, B. 1969. Effect of oat seed treatment with microelements. *Plant Soil* 31, 185-187.
59. Sabir Ahmed, A. (1989). Studies on production of quality seed and storage in soyabean. M.Sc. (Ag.) Thesis Tamil Nadu Agricultural University, Coimbatore.
60. Sneddon, J.L. Sugarbeet production experiments. *J. Natl. Inst. Agric. Bot.* 9 (1963): 333-345. according to Das (1990),
61. Scott. R.K. The sowing and harvesting dates, plant population and fertilizers on seed yield and quality of direct-drilled sugarbeet seed crops. *J. Agric. Sci.* 73 (1969): 373-385
62. Scofield GN, Aoki N, Hirose T, Takano M, Jenkins CL, Furbank RT (2007) The role of the sucrose transporter, OsSUT1, in germination and early seedling growth and development of rice plants. *J Exp Bot* 58:483-495. doi:10.1093/jxb/erl21
63. Singh, D. K., and S. Bharti. 1985. Seed manganese content and its relationship with the growth characteristics of wheat cultivar
64. Stout, P.R., Arnon, D.I. 1939. Experimental methods for the study of the role of copper, manganese, and zinc in the nutrition of higher plants. *Amer J. Bot.* 26, 144-149.
65. Saeed B, Gul H, Khan A.Z, Badshah N.L, Parveen L, Khan A. (2012). Rates and methods of nitrogen and sulfur application influence and cost benefit analysis of wheat. *Journal of Agricultural & Biological Science.* 7(2):81-85.
66. Sharma, A., H. T. Nakul, B. R. Jelgeri, and A. Surwenshi. 2010. Effect of mineral nutrients on growth, yield and yield components
67. in pigeonpea (*Cajanus cajan* L. Millsp.). *Research Journal of Agricultural Science* 1 (2):142-144.
68. Shibles R, Anderson TC, Gibson AH. 1975. Soybean. In: Evans LT, ed. *Crop physiology: some case histories.* Cambridge University Press, 151-189
69. Siegenthaler PA, Douet-Orhant V (1994) Relationship between the ATP content measured at three imbibition times and germination of onion seeds during storage at 3, 15 and 30 C. *J Exp Bot* 45:1365-1371. doi: 10.1093/jxb/45.10.1365
70. Singh, D. K., and S. Bharti. 1985. Seed manganese content and its relationship with the growth characteristics of wheat cultivars. *New Phytologist* 101:387-391.
71. Smil, V. 2000. Phosphorus in the environment: Natural flows and human interferences. *Annual Review of Energy and the Environment* 25: 53-88. doi:10.1146/annurev.energy.25.1.53
72. Shekhar C, Yadav A.L, Singh H.K, Singh M.K. (2010). Influence of micronutrients on plant growth, yield and quality of papaya fruit (*Carica papaya* L.) cv. Washington. *Asian J. Hort.* 5 (2): 326-329.
73. Sourour M.M. (2000). Effect of foliar application of some micronutrients forms on growth, yield, fruit quality and leaf mineral composition of Valencia orange trees grown in North Sinai. *Alexandria J. Agri. Res.* 45 (1): 269-285.
74. Stangoulis JCR, Grewal HS, Bell RW, Graham RD (2000) Boron efficiency in oilseed rape: I. Genotypic variation demonstrated in field and pot grown *Brassica napus* L. and *Brassica juncea* L. *Plant Soil* 225:243-251. doi:10.1023/a:1026593528256
75. Zagade P.M, Munde G.R, Shirsath A.H. (2017). Effect of foliar application of micronutrients on yield and quality of Guava (*Psidium guajava* L.) Cv. Sardar. *Journal of Pharmacy and Biological Sciences* 12(5), 56-58.
76. Talooth, A. T., M. H. Taha, and M. A. El-Seessy. 1989. Response of chemical composition and yield of soybean to foliar spray of Zn and different phosphatic fertilizers. *Annals of Agric. Sci. Cairo.* 34: 925-937.
77. TeKrony, Dennis M., and Dennis B. Egli. "Relationship of seed vigor to crop yield: a review." *Crop science* 31.3 (1991): 816-822.
78. Thompson LR, Torch FR. *Soils and soil fertility.* McGraw Hill Book Co. New York. 1978;226-283.
79. Vigidal SM, Pereira PRG, Pacheco DD. Mineral nutrition and fertilization of onion. *Informe. Agropecuario.* 2002;23:36-50.

80. Sing SP, Verma AB. Response of onion (*Allium cepa* L.) to potassium application. *Indian J. Agron.* 2001;46:182-185.
81. Yang M, Shi L, Xu F, Lu J, Wang Y 2009: Effects of B, Mo, Zn, and their interactions on seed yield of rapeseed (*Brassica napus* L.). *Pedosphere*, 19, 53–59. doi:10.1016/S1002-0160(08)60083-1
82. Wang T, Sistrunk LA, Leskovar DI, Cobb BG (2011) Characteristics of storage reserves of triploid watermelon seeds: association of starch and mean germination time. *Seed Sci Technol* 39: 318-326
83. Weitbrecht K, Müller K, Leubner-Metzger G (2011) First off the mark: early seed germination. *J Exp Bot* 62:3289-3309. doi: 10.1093/jxb/err030
84. Wedding RT, Black MK (1988) Role of magnesium in the binding of substrate and effectors to phosphoenolpyruvate carboxylase from a CAM plant. *Plant Physiol* 87:443-446. doi: 10.1104/pp.87.2.443
85. Wolswinkel P 1992 Transport of nutrients into developing seeds: a review of physiological mechanisms. *Seed Sci. Res.* 2
86. Zhang DD, Zhao H, Shi L, Xu FS (2014) Physiological and genetic responses to boron deficiency in *Brassica napus*: a review. *Soil Science and Plant Nutrition* 60:304-313. doi: 10.1080/00380768.2014.893537.
87. Zhao ZK, Wu LK, Nian FZ, Ding GD, Shi TX, Zhang DD, Shi L, Xu FS, Meng JL (2012) Dissecting quantitative trait loci for boron efficiency across multiple environments in *Brassica napus*. *PLoS One* 7:14. doi:10.1371/journal.pone.0045215